



Larimar Therapeutics

Nomlabofusp Program Update

December 2024

Forward-Looking Statements

This presentation contains forward-looking statements that are based on the beliefs and assumptions of Larimar Therapeutics, Inc. (“Company”) and on information currently available to management. All statements contained in this presentation other than statements of historical fact are forward-looking statements, including but not limited to Larimar’s ability to develop and commercialize nomlabofusp (CTI-1601) and other planned product candidates, Larimar’s planned research and development efforts, including the timing of its nomlabofusp clinical trials and non-clinical investigations and overall development plan expectations with respect to the FDA START pilot program, interactions with FDA, expectations regarding potential for accelerated approval or accelerated access and time to market and other matters regarding Larimar’s business strategies, ability to raise capital, use of capital, results of operations and financial position, and plans and objectives for future operations.

In some cases, you can identify forward-looking statements by the words “may,” “will,” “could,” “would,” “should,” “expect,” “intend,” “plan,” “anticipate,” “believe,” “estimate,” “predict,” “project,” “potential,” “continue,” “ongoing” or the negative of these terms or other comparable terminology, although not all forward-looking statements contain these words. These statements involve risks, uncertainties and other factors that may cause actual results, performance, or achievements to be materially different from the information expressed or implied by these forward-looking statements. These risks, uncertainties and other factors include, among others, the success, cost and timing of Larimar’s product development activities, nonclinical studies and clinical trials, including nomlabofusp clinical milestones and continued interactions with the FDA; that preliminary clinical trial results may differ from final clinical trial results, that earlier non-clinical and clinical data and testing of nomlabofusp may not be predictive of the results or success of later non-clinical or clinical trials, and assessments; that the FDA may not ultimately agree with Larimar’s nomlabofusp development strategy; the potential impact of public health crises on Larimar’s future clinical trials, manufacturing, regulatory, nonclinical study timelines and operations, and general economic conditions; Larimar’s ability and the ability of third-party manufacturers Larimar engages, to optimize and scale nomlabofusp’s manufacturing process; Larimar’s ability to obtain regulatory approvals for nomlabofusp and future product candidates; Larimar’s ability to develop sales and marketing capabilities, whether alone or with potential future collaborators, and to successfully commercialize any approved product candidates; Larimar’s ability to raise the necessary capital to conduct its product development activities; and other risks described in the filings made by Larimar with the Securities and Exchange Commission (SEC), including but not limited to Larimar’s periodic reports, including the annual report on Form 10-K, quarterly reports on Form 10-Q and current reports on Form 8-K, filed with or furnished to the SEC and available at www.sec.gov. These forward-looking statements are based on a combination of facts and factors currently known by Larimar and its projections of the future, about which it cannot be certain. As a result, the forward-looking statements may not prove to be accurate. The forward-looking statements in this presentation represent Larimar’s management’s views only as of the date hereof. Larimar undertakes no obligation to update any forward-looking statements for any reason, except as required by law.

Positive Initial Data from Long-Term OLE Study in FA

Daily 25 mg nomlabofusp administered in 14 participants for up to 260 days

Generally well-tolerated with long-term daily administration

- Generally well tolerated for over 8 months
- SAEs occurred in two study participants that resolved in 24 hours
- Most common AEs were mild injection site reactions

25 mg daily increased and maintained tissue FXN levels over time

- Tissue FXN levels showed mean change from baseline of 1.32 pg/ μ g in buccal cells and 9.28 pg/ μ g in skin cells at Day 90
- Increased from a mean of 15% of HV at baseline to 30% of HV in buccal cells and from 16% of HV to 72% of HV in skin cells at Day 90
- Tissue FXN levels appear to reach steady state levels by Day 30 in buccal cells

Predictable long-term pharmacokinetics

- Rapidly absorbed after subcutaneous administration
- Reached steady state levels in plasma by Day 30 with no further accumulation
- Pharmacokinetic profile consistent with Phase 1 and Phase 2 studies

Early trends of improvement observed in clinical outcomes at Day 90

- Early trends in mFARS, FARS-ADL, Modified Fatigue Impact Scale and 9 Hole Peg Test support the potential that daily nomlabofusp may lead to clinical benefit in patients with FA

Updates for Nomlabofusp Development Program

OLE Study

Six participants currently receiving 50 mg daily

Other enrolled participants will increase dose

All newly enrolled participants starting at 50 mg daily

Long-term 50 mg data expected mid-2025

Pediatric PK Run-In Study

Screening ongoing in adolescents (12-17 yrs)

Dosing expected to initiate in early 2025 at weight-based dose equivalent dose of adult 50 mg dose

Children (2-11 yrs) to begin enrollment in 1H 2025

Global Confirmatory/Registration Study

Identifying global sites in US, EU, UK, Australia and Canada

Finalizing protocol based on advice from global regulatory agencies

Initiation on track for mid-2025

BLA Submission/Accelerated Approval

Advancing discussions with FDA on data package required for FXN as a surrogate endpoint

Advancing discussions with FDA on the amount of the required safety data

BLA submission targeted for 2H 2025

Clinical-Stage Novel Protein Replacement Therapy Platform

Potential first therapy to systemically address route cause of FA

Lead candidate nomlabofusp is a recombinant fusion protein designed to directly address frataxin deficiency in patients with FA by delivering the protein to mitochondria. Granted Orphan Drug (US & EU), Rare Pediatric Disease (US), Fast Track (US), PRIME (EU) and ILAP (UK-MHRA) designations. Selected by FDA to participate in its START pilot program

Consistent Phase 1 and Phase 2 findings

Nomlabofusp was generally well tolerated and demonstrated dose-dependent increases in frataxin (FXN) levels from baseline in skin and buccal cells in completed studies

Advancing clinical program

Increased dose to 50 mg on current participants and starting newly enrolled participants on 50 mg in long-term OLE study with 50 mg data expected mid-2025
Screening of adolescents with FA ongoing in pediatric PK run-in study with dosing expected to begin in early 2025; adolescents completing study will transition into OLE after assessment of safety and PK data
Initiation of global confirmatory/registration study on track for mid-2025

Pursuing accelerated approval path with FDA

FDA acknowledgement that FXN deficiency appears to be critical to the pathogenic mechanism of FA, and that there continues to be an unmet need for treatments that address the underlying disease pathophysiology. Discussions to support an accelerated approval pathway are ongoing. BLA submission targeted for 2H 2025

Strong financial foundation

Approximately \$204 million in cash and investments as of 9/30/24, providing projected cash runway into Q2 2026

Friedreich's Ataxia (FA): A rare and progressive disease

Genetic defect on both alleles
lowers frataxin levels

Affects ~20,000
patients globally

Progressive disease

No approved therapies
increase frataxin levels

Most patients with FA only produce ~20-40% of normal frataxin levels depending on the tissue, sampling technique, and assay considered*

~5,000 patients in the U.S., with most remaining patients in the Europe
~70% of patients present before age 14

Initial symptoms include unsteady posture and frequent falling, and patients are eventually confined to a wheelchair

Life expectancy of 30-50 years with an early death usually caused by heart disease

Only treatment approved for FA does not address frataxin deficiency



FXN Levels Predict Disease Progression in FA

Lower FXN levels are associated with earlier onset of disease, faster rate of disease progression, and shorter time to loss of ambulation

Median Age of Onset and Rate of Disease Progression in Relation to FXN Levels

FXN Level* (% of Normal Level)	Age of Onset (Years)	FARS** (Change/Year)
11.2	7	2.9
22.0	11	2.1
31.0	16	2.0
48.7	19	1.6

Adapted from H.L.Plasterer et al. PLoS ONE 2013 8(5):e63958

Median Age of Onset Predicts Time to Loss of Ambulation

Age of Onset (Years)	Median Time to Loss of Ambulation (Years)
< 15	11.5
15 to 24	18.3
> 24	23.5

Adapted from C. Rummey et al. EClinicalMedicine. 2020 18:100213

Open-label Extension: 25 mg Completed, Increasing to 50 mg Daily

Key Eligibility Criteria

Previous participation in Phase 1 or Phase 2 trials

Daily subcutaneous injections self-administered or by a caregiver

25 mg nomlabofusp

50 mg nomlabofusp

- All 7 sites activated and enrolling
- Following assessment of safety and PK data of each cohort in the PK run-in study, adolescents (12-17 yrs) and children (2-11 yrs) will be eligible to screen for the OLE

Key Study Objectives

- Safety and tolerability
- Long-term PK
- Tissue FXN concentrations
- Clinical efficacy measures compared to FACOMS* database once enrollment is complete

Long-term data from 50 mg dose expected mid-2025

Screening Period \leq 42 days**


Treatment Period Planned for \geq 1 year

Potential extensions


Increased FXN Levels in Skin Cells Sustained Over Time

Participants dosed daily with 25 mg nomlabofusp for up to 90 days

Skin Cells - Absolute FXN Levels



Skin Cells - FXN Levels Change from Baseline




FXN levels measured via detection of peptide derived from mature FXN; FXN concentrations are normalized to total cellular protein content in each sample. Data represent median and 25th and 75th percentiles. Only participants with quantifiable levels at all measurement points are included in the figures. 50% of normal healthy volunteer (NHV) FXN level is 8.17pg/µg from the noninterventional healthy volunteer study (N=60).


Increased FXN Levels in Buccal Cells Sustained Over Time

Participants dosed daily with 25 mg nomlabofusp for up to 90 days and reached steady state by 30 days

Buccal Cells - Absolute FXN Levels





Buccal Cells - FXN Levels Change from Baseline



FXN levels measured via detection of peptide derived from mature FXN; FXN concentrations are normalized to total cellular protein content in each sample. Data represent median and 25th and 75th percentiles. Only participants with quantifiable levels at all measurement points are included in the figures. 50% of normal healthy volunteer (NHV) FXN level is 4.12 pg/µg from the noninterventional healthy volunteer study (N=60).

Tissue FXN Levels are Higher as a % of Healthy Volunteers at Day 90



Only participants with quantifiable levels at each measurement point are included in the tables
Mean % of healthy volunteers is the mean of all the participants FXN levels relative to the mean FXN levels in skin cells (16.34 pg/ μ g) and in buccal cells (8.24 pg/ μ g) from the noninterventional healthy volunteer study (N=60).

Observed Increases in Tissue FXN Levels in OLE Are Comparable to the Phase 2 Dose Exploration Study

Absolute tissue FXN levels and increases from baseline after 25 mg nomlabofusp daily over time

	Open Label Extension					
	Buccal FXN levels (pg/μg)			Skin FXN levels (pg/μg)		
	n	Median	Mean	n	Median	Mean
Baseline	11	1.13	1.19	8	2.41	2.60
Day 30	11	2.08	3.62	8	5.34	7.45
Change from Baseline	11	0.58	2.43	8	2.42	4.85
Day 60	9	2.46	2.41			
Change from Baseline	9	0.53	1.13			
Day 90	6	1.89	2.48	5	7.65	11.73
Change from Baseline	6	1.01	1.32	5	4.89	9.28

	Phase 2 Dose Exploration					
	Buccal FXN levels (pg/μg)			Skin FXN levels (pg/μg)		
	n	Median	Mean	n	Median	Mean
Baseline	7	1.78	1.80	7	3.70	3.38
Day 14	7	2.24	2.22	7	5.53	6.4
Change from Baseline	7	0.56	0.42	7	2.81	3.02

Observed Trends Towards Improvement in Clinical Outcomes at Day 90 After Daily 25 mg Nomlabofusp

Visit	Statistic	mFARS 93-Point Scale	FARS-ADL 36-Point Scale	Modified Fatigue Impact Scale 84-Point Scale	9 Hole Peg Test Dominant Hand Time (Seconds)
		N = 8	N = 8	N = 8	N = 8
Baseline	Mean (SD)	55.81 (13.296)	18.13 (6.064)	27.1 (14.23)	130.91 (99.366)
	Median (IQR)	53.5 (47.5, 68.3)	17.0 (12.8, 23.8)	29.5 (18, 38)	89.5 (48.7, 227.8)
	(Min, Max)	(35.0, 73.0)	(11.0, 27.0)	(2, 45)	(38.0, 277.3)
Day 90	Mean (SD)	55.13 (14.829)	15.88 (6.249)	18.5 (15.68)	113.11 (95.586)
	Median (IQR)	53.3 (43.8, 66.0)	14.8 (11.0, 21.3)	17.0 (5, 32)	67.15 (48.4, 176.7)
	(Min, Max)	(35.3, 79.5)	(8.0, 25.0)	(0, 42)	(33.50, 287.00)
Change from Baseline at Day 90	Mean (SD)	-0.69 (3.983)	-2.25 (3.082)	-8.6 (12.24)	-17.79 (27.450)
	Median (IQR)	-1.17 (-3.8, 1.2)	-2.25 (-3.8, 0.3)	-3.5 (-19, -3)	-9.00 (-32.0, 1.7)
	(Min, Max)	(-5.0, 7.0)	(-8.0, 1.5)	(-28, 9)	(-73.5, 9.8)

Nomlabofusp: Predictable Long-Term Pharmacokinetics

1

Rapid absorption after subcutaneous administration

2

Nomlabofusp reached steady state in plasma by Day 30 with no further accumulation

3

Pharmacokinetic profile consistent with Phase 1 and Phase 2 studies

OLE Study: Nomlabofusp 25 mg Daily is Generally Well-Tolerated for Up to 260 Days

14 participants with FA were included in the safety data

One additional participant started dosing after the data cut and one additional participant is scheduled for dosing before year-end 2024

Two participants withdrew for non-treatment related reasons

Two participants had serious adverse events that resolved within 24 hours and withdrew from the study

- Events were reviewed by Data Monitoring Committee and submitted to FDA and study is continuing as planned

Most common adverse events (AEs) were injection site reactions (ISRs) which were mild, brief in duration and self limited

No study discontinuations due to ISRs and all resolved

Increasing nomlabofusp dose to 50 mg in participants already enrolled and starting newly enrolled participants on 50 mg daily

Totality of Data to Date Supports that Nomlabofusp has the Potential to Slow Disease Progression

Addresses underlying deficiency

FDA alignment that FXN deficiency appears to be the pathogenic mechanism of FA

Dose dependent increases in FXN levels that were maintained over time

Displays functional activity

Trends in modified gene expression and lipid profile changes towards values seen in healthy controls

Data suggest nomlabofusp may positively affect downstream metabolic pathways disrupted in FA


Early trend in clinical benefit

Trends towards improvement observed in multiple clinical outcome measures

Potential to benefit ambulatory and non-ambulatory patients with FA

Nomlabofusp Clinical Development Program

Intend to pursue accelerated approval pathway with potential BLA submission targeted for 2H 2025
Selected by FDA to participate in its START pilot program



Ongoing open-label extension study with 50 mg daily dosing for eligible patients who participated in Phase 1 and/or dose exploration studies

Long-term data from 50 mg dose expected mid-2025



Pediatric PK run-in study initiated in adolescents (12-17 yrs), followed by children (2-11 yrs) in 1H 2025

Participants completing the PK run-in study eligible to transition into OLE after assessment of safety and PK data from each cohort



Planned global double-blind placebo-controlled confirmatory/registration study on track to initiate mid- 2025*

BLA submission targeted for 2H 2025**

Positive Long-term OLE Data and Advancing Dose to 50 mg

OLE Supports Potential of Nomlabofusp

Nomlabofusp was generally well tolerated at doses administered up to 260 days

Tissue FXN levels showed mean change from baseline of 1.32 pg/μg in buccal cells and 9.28 pg/μg in skin cells at Day 90

Daily nomlabofusp 25 mg increased and maintained tissue FXN over time, increasing from a mean of 15% of HV at baseline to 30% of HV at Day 90 in buccal cells and from 16% of HV at baseline to 72% of HV at Day 90 in skin cells

Early trends towards improvements across multiple clinical outcomes at Day 90, supporting potential clinical benefit

Clinical & Regulatory Updates

Increasing dose to 50 mg in OLE study with 6 participants administered 50 mg to date

Screening adolescents for pediatric PK run-in study

Evaluating global clinical sites for planned registration/confirmatory study

Selected by FDA to participate in its START pilot program

Ongoing discussions with FDA regarding use of FXN as a surrogate endpoint to support accelerated approval pathway

2025 Milestones

Q1 2025: Dose adolescents (ages 12-17 years old) in pediatric run-in study

1H 2025: Initiate PK run-in study in children (ages 2-11 years old)

Mid 2025: Initiate global confirmatory/registration study

Mid 2025: Data from 50 mg dose in OLE study

2H 2025: BLA submission; intend to pursue accelerated approval



Larimar Therapeutics


Appendix

Larimar Technology is Supported by a Strong IP Portfolio


Granted nomlabofusp (CTI-1601) composition of matter patent extends into 2040

2010 2015 2020 2025 2030 2035 2040 2045 2050 2055

Composition of Matter Patent



Formulation and Methods of Quantifying Nomlabofusp



Platform Applications



Additional nomlabofusp IP protection

- US and foreign pending applications cover key biomarkers, analytical tools and methods of treatment for additional disease indications for nomlabofusp
- Nomlabofusp should be eligible for **12 years of market exclusivity** upon approval in the US (independent of patents) and at least **10 years of market exclusivity** upon approval in EU (independent of patents)

Nomlabofusp is Designed to Deliver Additional Frataxin

Nomlabofusp (CTI-1601) maintains the cleavage site between the MTS and mature human frataxin (FXN)

STRUCTURE OF ENDOGENOUS FXN

Mitochondrial Targeting Sequence (MTS)

Mature Human FXN



Cleavage by mitochondrial processing peptidase (MPP) at this site produces mature human FXN in mitochondria

STRUCTURE OF NOMLABOFUSP

Mitochondrial Targeting Sequence (MTS)

Mature Human FXN



Cell Penetrating Peptide (CPP)

Cleavage by mitochondrial processing peptidase (MPP) at this site produces mature human FXN in mitochondria

The presence of the cleavage site allows the CPP and MTS to be removed by mitochondrial processing peptidase to produce mature human FXN in the mitochondria

Nomlabofusp is a Competitively Differentiated Treatment Approach*

\$7.3B

Acquisition supports the **robust market potential** for FA treatments



Nomlabofusp is a potential **first-and-only protein replacement therapy** designed to address the underlying cause of FA


Approach	Product	Company	Mechanism of Action	Clinical Status
Protein replacement	Nomlabofusp (CTI-1601)	Larimar	Frataxin Protein Replacement	Phase II
Mitochondrial Oxidative Stress Modifier	Omaveloxolone (SKYCLARYS™)	Biogen	Nrf2 Activator	Approved (US and EU)
	Vatiquinone	PTC Therapeutics	15-Lipoxygenase Inhibitor	Phase III
Gene Expression Regulator	DT-216P2 (new formulation)	Design Therapeutics	GeneTAC	Pre-clinical
Gene Therapy	LX2006	Lexeo Therapeutics	Frataxin Gene Replacement	Phase I/II




Mitochondrial Localization and Preclinical Data

Nomlabofusp Transduction of Cells In Vitro Leads to hFXN Located in Mitochondria


FXN staining



TOMM20 (mitochondria) staining



FXN co-localizes with TOMM20



- Rat cardiomyocytes (H9C2) were transduced with nomlabofusp
- Cells were fixed and analyzed by immunofluorescence microscopy to detect the presence of human frataxin (hFXN) and TOMM20 (a mitochondrial outer membrane protein)
- Nuclei were stained with DAPI

Nomlabofusp Extends Survival in FXN-deficient KO Mice


Initial proof-of-concept for FXN replacement therapy in cardiac mouse model of FA

Median survival of MCK-Cre FXN-KO mice

- 166 days (nomlabofusp) vs. 98 days (Vehicle)
- Nomlabofusp administered 10 mg/kg SC every other day

Survival beyond vehicle mean (107.5 days)

- 87.5% (nomlabofusp) vs. 33% (Vehicle)
- Demonstrates that nomlabofusp is capable of delivering sufficient amounts of FXN to mitochondria



Nomlabofusp (CTI-1601) rescues a severe disease phenotype in a well-characterized cardiac mouse model of FA

Nomlabofusp Prevents Development of Ataxic Gait in Neurologic KO Mouse Model

In-Vivo Efficacy Data in Pvalb-Cre FXN-KO Mouse Model


Single dose level: 10 mg/kg nomlabofusp or vehicle given intraperitoneally three times per week

- ✓ hFXN replacement with nomlabofusp **prevents development of ataxic gait**
- ✓ Nomlabofusp-treated mice **survive longer** than untreated mice
- ✓ Human frataxin **present in brain, dorsal root ganglia and spinal cord** demonstrating central nervous system penetration

Nomlabofusp Delivers hFXN to Mitochondria and Restores SDH Activity in KO Mice


Study Design – Cardiac and skeletal muscle FXN knockout mice (MCK-CRE) were treated at varying SQ doses of nomlabofusp every other day for two weeks at Jackson Laboratories (Bar Harbor, ME). After dosing, animals were sacrificed, and heart and skeletal muscle were evaluated for hFXN concentration in mitochondrial extracts and SDH activity was assessed.

Mitochondrial FXN (Heart)



Mitochondria hFXN concentration increases dose-dependently
Given subcutaneously, nomlabofusp functionally replaces hFXN
in mitochondria of KO mice

SDH Activity (Muscle)




Succinate dehydrogenase (SDH) activity, which is indicative of mitochondrial function, increases in a dose-dependent manner after administration of nomlabofusp; activity plateaus at 30 mg/kg and is equivalent to activity in wild type


Nomlabofusp Prevents Left Ventricle Dilatation in KO Mice

Study Design – Cardiac and skeletal muscle FXN knockout mice (MCK-CRE) were treated at 10 mg/kg every other day at Jackson Laboratories (Bar Harbor, ME). Echocardiograms were performed pre-dose and post dose.

Left Ventricle Internal Diameter (Systole)



Left Ventricle Volume (Systole)




Left ventricular (LV) volume increases in systole in untreated mice by 8 weeks (after 4 weeks of dosing with vehicle), but remains similar to wildtype when treated with nomlabofusp (10 mg/kg every other day)

Nomlabofusp-treated mice have similar LV volume as wild type; echocardiogram shows significant differences between vehicle and nomlabofusp treated (10 mg/kg every other day) KO mice


Nomlabofusp Preserves Left Ventricle Function in KO Mice

Study Design – Cardiac and skeletal muscle FXN knockout mice (MCK-CRE) were treated at 10 mg/kg every other day at Jackson Laboratories (Bar Harbor, ME). Echocardiograms were performed pre-dose and post dose.

Left Ventricle Ejection Function



Left Ventricle Fractional Shortening



Left ventricular (LV) function drops significantly in vehicle treated mice by Week 8

Nomlabofusp-treated (10 mg/kg every other day) mice have similar LV function as wildtype; echocardiogram shows significant differences between vehicle and nomlabofusp treated KO mice




Phase 1 Clinical Data

CTI-1601: Phase 1 Clinical Program in Patients with FA

Program consisted of double-blind, placebo controlled single- and multiple-ascending dose trials

Phase 1 Development Plan

- Two double-blind, placebo-controlled dosing trials in patients with FA
- Patient dosing began December 2019
- Safety Review Committee assessed all blinded data between each cohort to ensure patient safety



Number of subjects: 28

Dose levels: 25 mg, 50 mg, 75 mg and 100 mg (subcutaneous administration)

Treatment Duration: 1 day

1^o Endpoint: Safety and tolerability

2^o Endpoints: PK; PD; FXN levels; multiple exploratory

Status: Complete

Number of Subjects: 27

Dose Range: 25 mg, 50 mg, 100 mg (subcutaneous administration)

Treatment Regimen: Multiple increasing doses administered subcutaneously over 13 days




1^o Endpoint: Safety and tolerability

2^o Endpoints: PK; PD; FXN levels (buccal cells, platelets, optional skin biopsies); multiple exploratory

Status: Complete

Completed Phase 1 Multiple Ascending Dose Study

Treatment Schedules for Each Cohort- nomlabofusp (CTI-1601) or placebo



FXN Level Sampling Days Presented for Each Cohort


Cohort 1 Sampling Days	
Buccal Cells	Baseline, Day 4, Day 13
Skin	Baseline, Day 13
Platelets	Baseline, Day 4, Day 13

Cohort 2 Sampling Days	
Buccal Cells	Baseline, Day 7, Day 13
Skin	Baseline, Day 13
Platelets	Baseline, Day 7, Day 13

Cohort 3 Sampling Days	
Buccal Cells	Baseline, Day 7, Day 13
Skin	Baseline, Day 13
Platelets	Baseline, Day 7, Day 13


Dose Dependent Increases in FXN Levels Observed in Skin and Buccal Cells in Phase 1

FXN* Change from Baseline By Dose Group (Skin Cells)



Placebo: Participants randomized to placebo in each cohort
25 mg: Dosed daily for 4 days, every third day thereafter

FXN* Change from Baseline By Dose Group (Buccal Cells)



50 mg: Dosed daily for 7 days, every other day thereafter
100 mg: Dosed daily for 13 days

*FXN levels measured via detection of peptide derived from mature FXN; FXN concentrations are normalized to total cellular protein content in each sample; Data represent median and 25th and 75th percentiles; FXN levels from Day 4, & Day 13 measurements are shown for data derived from the 25 mg cohort; FXN levels from Day 7 & Day 13 measurements are shown for data derived from the 50 & 100 mg cohorts;

MAD Trial Patient Demographics

Parameter	Statistic	All placebo (n=7)	25 mg CTI-1601 (n=6)	50 mg CTI-1601 (n=7)	100 mg CTI-1601 (n=7)	All CTI-1601 (n=20)	Overall (n=27)
Sex							
Male	n (%)	5 (71.4)	3 (50.0)	4 (57.1)	3 (42.9)	10 (50.0)	15 (55.6)
Female	n (%)	2 (28.6)	3 (50.0)	3 (42.9)	4 (57.1)	10 (50.0)	12 (44.4)
Age (years)							
	Mean	25.7	39.7	34.7	28.0	33.9	31.7
	SD	6.37	16.59	9.03	8.96	12.13	11.40
	Median	23	37	36	24	34	28
	Min, Max	20,36	21,65	19,47	20,44	19,65	19,65
Race							
White	n (%)	6 (85.7)	6 (100.0)	6 (85.7)	6 (85.7)	18 (90.0)	24 (88.9)
Asian	n (%)	0	0	1 (14.3)	1 (14.3)	2 (10.0)	2 (7.4)
American Indian	n (%)	1 (14.3)	0	0	0	0	1 (3.7)
Ethnicity							
Hispanic/Latino	n (%)	2 (28.6)	0	0	0	0	2 (7.4)
Not Hispanic/Latino	n (%)	5 (71.4)	6 (100.0)	7 (100.0)	7 (100.0)	20 (100.0)	25 (92.6)

MAD Trial Patient Disease Characteristics

Parameter	Statistic	All placebo (n=7)	25 mg CTI-1601 (n=6)	50 mg CTI-1601 (n=7)	100 mg CTI-1601 (n=7)	All CTI-1601 (n=20)	Overall (n=27)
Age at Symptom Onset							
	Mean	14.1	24.0	19.3	11.9	18.1	17.1
	SD	5.34	14.48	6.21	6.72	10.37	9.39
	Median	15.0	18.0	19.0	10.0	18.0	16.0
	Min, Max	8,23	12,44	8,28	5,22	5,44	5,44
Age at Diagnosis							
	Mean	18.3	31.5	26.4	15.9	24.3	22.7
	SD	7.87	19.88	4.28	8.21	13.24	12.23
	Median	20.0	25.5	28.0	13.0	27.0	21.0
	Min, Max	9,32	14,64	17,30	5,27	5,64	5,64
Assistive Device							
Walker	n (%)	0	2 (33.3)	3 (42.9)	0	5 (25.0)	5 (18.5)
Wheelchair	n (%)	4 (57.1)	3 (50.0)	1 (14.3)	6 (85.7)	10 (50.0)	14 (51.9)
Other	n (%)	1 (14.3)	0	1(14.3)	0	1 (5.0)	2 (7.4)
None	n (%)	2 (28.6)	1 (16.7)	2 (28.6)	1 (14.3)	4 (20.0)	6 (22.2)

PK analyses support evaluating once-daily and every-other-day dosing regimens for CTI-1601

Summary of MAD Trial PK Analyses

- ✓ CTI-1601 was quickly absorbed after subcutaneous administration
- ✓ Dose-proportional increases in exposure observed with increasing doses of CTI-1601
- ✓ Mean half life of CTI-1601 in plasma was approximately 11 hours
- ✓ CTI-1601 appeared to be at or close to steady state exposure after 13 days of dosing 100 mg once daily



Phase 2 Dose Exploration Data

Completed Ph 2 Dose Exploration Study (25 & 50 mg Cohorts)

Goal: Further characterize PK/PD and assess safety to inform long-term dose and dose regimen

28-day Treatment Period - nomlabofusp (CTI-1601) or placebo



■ = Subcutaneous administration of nomlabofusp (CTI-1601) or placebo

■ = No Administration


Study Details

Population	Ambulatory and non-ambulatory Friedreich's ataxia patients ≥ 18 years of age Nomlabofusp (CTI-1601) treatment naïve or participated (if eligible) in a previous Larimar study
Dose	Cohort 1: 25 mg Cohort 2: 50 mg
Key Endpoints	Frataxin levels in peripheral tissue, PK, safety and tolerability; other exploratory endpoints include lipids and gene expression levels
Number of Patients	Cohort 1: Enrolled 13 participants (9 on nomlabofusp; 4 on placebo) Cohort 2: Enrolled 15 participants (10 on nomlabofusp; 5 on placebo)
Key Results	Generally well tolerated; most common adverse events were mild and moderate injection site reactions Dose dependent increases of frataxin levels in tissues tested (skin and buccal cells) Baseline FXN levels in skin cells in the 50 mg cohort were $< 17\%$ of the average of healthy volunteers. After daily dosing for 14 days, FXN levels increased to 33% to 59% of the average of the healthy volunteers


Dose-Dependent Increase in FXN Levels in Skin Cells

Participants dosed daily for 14 days, then every other day until day 28

Skin Cells FXN Levels* Change from Baseline**



FXN Levels* in Skin Cells Change from Baseline at Day 14




*FXN levels measured via detection of peptide derived from mature FXN; FXN concentrations are normalized to total cellular protein content in each sample. Data represent median and 25th and 75th percentiles. Only participants with quantifiable levels at both baseline and Day 14 are included in the figures.

**Median baseline FXN levels in patients were 3.5 pg/μg for the placebo, 3.7 pg/μg for the 25 mg cohort and 2.1 pg/μg for the 50 mg cohort.


Dose-Dependent Increase in FXN Levels in Buccal Cells

Participants dosed daily for 14 days, then every other day until day 28

Buccal Cells FXN Levels* Change from Baseline**



FXN Levels* in Buccal Cells Change from Baseline at Day 14




*FXN levels measured via detection of peptide derived from mature FXN; FXN concentrations are normalized to total cellular protein content in each sample. Data represent median and 25th and 75th percentiles. Only participants with quantifiable levels at both baseline and Day 14 are included in the figures.


**Median baseline FXN level in patients were 2.1 pg/μg for the placebo, 1.8 pg/μg for the 25 mg cohort and 1.6 pg/μg for the 50 mg cohort.

Skin Cell FXN Levels Achieve Higher % of Healthy Volunteers* Following 14 days of Daily Nomlabofusp

25 mg of Nomlabofusp



50 mg of Nomlabofusp




■ Baseline FXN levels as a % of average FXN level in healthy volunteers

■ FXN levels increased from baseline and reached 25% to < 50% of average FXN level in healthy volunteers


■ FXN levels increased from baseline and reached > 50% of average FXN level in healthy volunteers

Buccal Cell FXN Levels Achieve Higher % of Healthy Volunteers* Following 14 days of Daily Nomlabofusp

25 mg of Nomlabofusp



50 mg of Nomlabofusp



■ Baseline FXN levels as a % of average FXN level in healthy volunteers

■ FXN levels increased from baseline and reached 25% to < 50% of average FXN level in healthy volunteers

Absolute Increases in Skin FXN Levels

Dose response in tissue FXN concentrations and increases from baseline after dosing

Day 14 Skin FXN Levels			
Dose	Visit	Absolute Values (pg/μg)	
		Median	Mean
25 mg	Baseline	3.70	3.38
	Day 14	5.53	6.40
	Change from Baseline	2.81	3.02
50 mg	Baseline	2.12	2.08
	Day 14	7.40	7.32
	Change from Baseline	5.57	5.24

Day 28 Skin FXN Levels			
Dose	Visit	Absolute Values (pg/μg)	
		Median	Mean
25 mg	Baseline	3.70	3.38
	Day 28	4.39	4.80
	Change from Baseline	2.28	1.41
50 mg	Baseline	2.12	2.08
	Day 28	5.23	5.24
	Change from Baseline	3.14	3.17

Absolute Increases in Buccal FXN Levels

Dose response in tissue FXN concentrations and increases from baseline after dosing

Day 14 Buccal FXN Levels			
Dose	Visit	Absolute Values (pg/μg)	
		Median	Mean
25 mg	Baseline	1.78	1.80
	Day 14	2.24	2.22
	Change from Baseline	0.56	0.42
50 mg	Baseline	1.61	1.69
	Day 14	2.44	2.38
	Change from Baseline	0.72	0.69

Day 28 Buccal FXN Levels			
Dose	Visit	Absolute Values (pg/μg)	
		Median	Mean
25 mg	Baseline	1.70	1.65
	Day 28	1.73	1.76
	Change from Baseline	0.03	0.11
50 mg	Baseline	1.76	1.77
	Day 28	2.15	2.15
	Change from Baseline	0.48	0.38

Nomlabofusp: Predictable Pharmacokinetics

1

Quick absorption after subcutaneous administration

2

Dose-proportional increases in exposure observed

3


Pharmacokinetic profile consistent with Phase 1 studies ^{CC}

MAD Trial Patient Demographics

Parameter	Statistic	All placebo (n=7)	25 mg CTI-1601 (n=6)	50 mg CTI-1601 (n=7)	100 mg CTI-1601 (n=7)	All CTI-1601 (n=20)	Overall (n=27)
Sex							
Male	n (%)	5 (71.4)	3 (50.0)	4 (57.1)	3 (42.9)	10 (50.0)	15 (55.6)
Female	n (%)	2 (28.6)	3 (50.0)	3 (42.9)	4 (57.1)	10 (50.0)	12 (44.4)
Age (years)							
	Mean	25.7	39.7	34.7	28.0	33.9	31.7
	SD	6.37	16.59	9.03	8.96	12.13	11.40
	Median	23	37	36	24	34	28
	Min, Max	20,36	21,65	19,47	20,44	19,65	19,65
Race							
White	n (%)	6 (85.7)	6 (100.0)	6 (85.7)	6 (85.7)	18 (90.0)	24 (88.9)
Asian	n (%)	0	0	1 (14.3)	1 (14.3)	2 (10.0)	2 (7.4)
American Indian	n (%)	1 (14.3)	0	0	0	0	1 (3.7)
Ethnicity							
Hispanic/Latino	n (%)	2 (28.6)	0	0	0	0	2 (7.4)
Not Hispanic/Latino	n (%)	5 (71.4)	6 (100.0)	7 (100.0)	7 (100.0)	20 (100.0)	25 (92.6)

MAD Trial Patient Disease Characteristics

Parameter	Statistic	All placebo (n=7)	25 mg CTI-1601 (n=6)	50 mg CTI-1601 (n=7)	100 mg CTI-1601 (n=7)	All CTI-1601 (n=20)	Overall (n=27)
Age at Symptom Onset							
	Mean	14.1	24.0	19.3	11.9	18.1	17.1
	SD	5.34	14.48	6.21	6.72	10.37	9.39
	Median	15.0	18.0	19.0	10.0	18.0	16.0
	Min, Max	8,23	12,44	8,28	5,22	5,44	5,44
Age at Diagnosis							
	Mean	18.3	31.5	26.4	15.9	24.3	22.7
	SD	7.87	19.88	4.28	8.21	13.24	12.23
	Median	20.0	25.5	28.0	13.0	27.0	21.0
	Min, Max	9,32	14,64	17,30	5,27	5,64	5,64
Assistive Device							
Walker	n (%)	0	2 (33.3)	3 (42.9)	0	5 (25.0)	5 (18.5)
Wheelchair	n (%)	4 (57.1)	3 (50.0)	1 (14.3)	6 (85.7)	10 (50.0)	14 (51.9)
Other	n (%)	1 (14.3)	0	1(14.3)	0	1 (5.0)	2 (7.4)
None	n (%)	2 (28.6)	1 (16.7)	2 (28.6)	1 (14.3)	4 (20.0)	6 (22.2)



*Data Presented at the
International Congress for
Ataxia Research, November
2024*

Nomlabofusp Clinical Studies Included a Broad, Representative Population of Adults with FA

Broad population of adults with FA included in Phase 1 and 2 Studies

Age of onset between 5 - 60 years with a median age of onset of 15 yrs

81% of participants had FXN levels at baseline less than 30% of healthy controls and 37% of participants had less than 20%

Over 50% of participants were non-ambulatory at baseline

**18 subjects participated in more than 1 study*

***Quantifiable buccal cell FXN levels relative to the median of healthy controls*

****Ambulatory status is based on the gait score (E7=5 vs. <5) of the upright stability subscore of the mFARS*

*****Data presented at the International Congress for Ataxia Research, November 2024*

Demographics and Baseline Disease Characteristics from Nomlabofusp Phase 1 and 2 Interventional Studies****

	N*	Median	Mean	Min	Max
Age	61	28.0	31.9	19	69
Age of Onset	61	15.0	15.9	5	60
Age of Diagnosis	61	19.0	21.0	5	64
Shorter GAA (GAA₁)	60	550.0	555.8	99	1000
Longer GAA (GAA₂)	60	900.0	890.2	265	1300
Frataxin, % of Control**	57	24.4	23.9	8.7	61.9
mFARS Score	61	52.0	49.5	13.2	74.5
Upright Stability Score	61	32.0	26.9	7.0	35.0
Dominant hand 9-hole peg test	61	71.0	84.8	26.0	229.2
T25-FW Test Score	51	9.9	13.4	4.3	48.5
Left Ventricular Mass (g)	61	163.4	168.0	73.7	398.8
LVEF %	61	63.0	63.5	52	76
Ambulatory Status***					
No	36				
Yes	25				

Pooled Data from Completed Phase 1 & 2 Studies Confirms Disease & FXN Relationships are Consistent with Literature

Disease Characteristics by Quartiles Based on Buccal Cell FXN Levels at Baseline

Quartile	FXN Concentration* (pg/mcg)	Age at Symptom Onset**	Age at Diagnosis**	GAA ₁ **	GAA ₂ **
Q1 (N=14)	< 1.31	10.5	14.5	616.5	899.5
Q2 (N=14)	1.31 - <1.95	13.5	23.0	486.0	866.0
Q3 (N=14)	1.95 - <2.30	16.0	19.0	555.0	871.5
Q4 (N=15)	≥ 2.30	19.0	27.0	400.0	933.0


*Quantifiable buccal cell frataxin levels

**Median values

Median buccal cell FXN concentration in healthy controls = 8.1 ng/mcg


Buccal cell FXN levels correlated with age of onset and inversely correlated with the number of GAA repeats and rate of disease progression

Baseline Buccal and Skin Cell FXN Levels



Buccal cell FXN levels correlated with skin cell FXN levels

Modeling/Simulation Predicts* 50mg Daily Can Achieve Skin FXN Levels $\geq 50\%$ of Healthy Controls in Most Patients



Dashed red line – 50% the average skin FXN/protein ratio (8.17 pg/ug) in a non-interventional study in healthy controls (HC)

Blue line – median of simulated values across trials

Red lines – 10th and 90th percentiles

Shaded regions – 95% confidence intervals of the corresponding percentiles (10th, 50th, and 90th).

Data presented at the International Congress for Ataxia Research, November 2024


50 mg nomlabofusp daily was predicted to lead to:

A median increase of 5.64 (2.3 – 13.5) pg/ μ g in FXN levels from baseline


Increase in skin FXN levels in 59% of simulated patients with FA to levels $\geq 50\%$ of average skin FXN levels in HC

Increase Towards Normal Gene Expression in Adults with FA* Observed After Nomlabofusp Treatment

Select Baseline Gene Expression Patients with FA* vs. Healthy Volunteers (HV)**



Post-treatment Changes in Gene Expression From Baseline




Data presented at the International Congress for Ataxia Research, November 2024

*Samples from Phase 2 dose exploration study evaluating nomlabofusp 25 mg (Cohort 1) and 50 mg (Cohort 2) or placebo via subcutaneous injection daily for 14 days followed by alternate day administration for 14 days. Buccal samples were collected before, during, and after treatment for gene expression profiling

**Data from Larimar's non-interventional healthy volunteer study

Decrease Towards Normal Lipid Profiles in Adults with FA* Observed After Nomlabofusp Treatment



*Samples from Phase 2 dose exploration study evaluating nomlabofusp 25 mg (Cohort 1) and 50 mg (Cohort 2) or placebo via subcutaneous injection daily for 14 days followed by alternate day administration for 14 days. Plasma samples were collected before, during, and after treatment for lipid profiling. Healthy volunteer (HV) data is from Larimar's non-interventional HV study

Data presented at the International Congress for Ataxia Research, November 2024




Non-Interventional Study Data

CLIN-1601-002: Top-line Non-interventional Study Results

Non-interventional study measured FXN in homozygous healthy volunteers

Median Frataxin Concentration (pg/ μ g)
in Homozygous Healthy Volunteers (n = 60)



Most patients with FA only produce ~20-40%¹ of normal frataxin levels depending on the tissue, sampling technique, and assay considered

Lower FXN levels seen with typical onset² (5 to 15 years of age)

Higher FXN levels seen with late onset² (after 25 years of age)

Heterozygous carriers who show no signs of disease have buccal cell FXN levels of ~50% of unaffected healthy persons¹



FDA START Pilot Program

Nomlabofusp Selected by FDA for START Pilot Program

Highlights FDA commitment to augment formal meetings with more rapid, ad-hoc communications to accelerate program development of rare diseases

START Pilot Program

Support for Clinical Trials Advancing Rare Disease Therapeutics

A new milestone-driven program launched by the FDA in September 2023

Designed to **accelerate development of novel therapies** intended to address unmet medical needs in **rare diseases**

7 novel drugs selected

- 3 products by CDER (nomlabofusp) for rare neurodegenerative conditions
- 4 products by CBER for cell and gene therapy

CDER Selection Based On

Demonstrated development **program readiness** (e.g., sponsors who demonstrate the ability to move the program towards a marketing application)

Potential to address serious and unmet medical need in a **rare neurodegenerative condition**

Alignment of CMC development timelines with clinical development plans

Proposed plan where **enhanced communication can improve efficiency of product development**



FARA

Strong Relationship with FARA – Joined FARA’s TRACK-FA Neuroimaging Consortium as an Industry Partner

TRACK-FA collects natural history data to establish disease specific neuroimaging biomarkers for potential use in clinical trials. Larimar will have access to all study data for use in regulatory filings, as appropriate

FARA provides industry with several key items

- Assistance with patient recruitment and education
- Access to Global Patient Registry with demographic and clinical information on more than 1,000 FA patients
- Sponsored a Patient-Focused Drug Development Meeting in 2017 resulting in a publication titled “The Voice of the Patient”



National, non-profit organization dedicated to the pursuit of scientific research leading to treatments and a cure for FA